

Chapter 1 Matter and Measurement

Solutions to Practice Problems

1.1 Representation (a) is a pure substance because each particle contains one red and two gray spheres. Representation (b) is a mixture because some of the particles are only red, and some are red and black.

1.2 Use Table 1.2 to determine the prefix for each unit.

- a. a million liters = megaliter
b. a thousandth of a second = millisecond
c. a hundredth of a gram = centigram
d. a tenth of a liter = deciliter

1.3 All nonzero digits are significant. A zero is significant only if it occurs between two nonzero digits, or at the end of a number with a decimal point. The significant figures are in **bold**.

- | | | | | |
|---|--|---|--|--|
| a. 23.45 4 significant figures | c. 230 2 significant figures | e. 0.202 3 significant figures | g. 1,245,006 7 significant figures | i. 10,040 4 significant figures |
| b. 23.057 5 significant figures | d. 231.0 4 significant figures | f. 0.003 60 3 significant figures | h. 1,200,000 2 significant figures | j. 10,040. 5 significant figures |

1.4 When the number to be rounded off is 4 or fewer, it and all other digits to the right are dropped. When the number is 5 or greater, 1 is added to the digit to its left.

- a. 1.2735 → 1.3
Since this number is 7 (5 or greater), round the 2 to its left up by one.
- b. 0.002 536 22 → 0.0025
Since this number is 3 (4 or fewer), drop it and all numbers to its right.
- c. 3,836.9 → 3,800
Since this number is 3 (4 or fewer), drop it and all numbers to its right.

1.5 The answers must have the same number of significant figures as the original number with the fewest number of significant figures.

a. $10.70 \times 3.5 = 37.45$
Because 3.5 has only two significant figures, round the answer to give it two significant figures.

37

c. $1,300 \div 41.2 = 31.553\ 398$
Because 1,300 has only two significant figures, round the answer to give it two significant figures.

32

b. $0.206 \div 25,993 = 0.000\ 007\ 93$

Because 0.206 has three significant figures, the answer has the appropriate number of significant figures.

d. $120.5 \times 26 = 3,133$

Because 26 has only two significant figures, round the answer to give it two significant figures.

3,100

1.6 The answers must have the same number of decimal places as the original number with the fewest decimal places.

a. $27.8\text{ cm} + 0.246\text{ cm} = 28.046\text{ cm}$

Because 27.8 has one digit after the decimal point, round the answer to one digit after the decimal point.

28.0 cm

b. $102.66\text{ mL} + 0.857\text{ mL} + 24.0\text{ mL} = 127.517\text{ mL}$

Because 24.0 has one digit after the decimal point, round the answer to one digit after the decimal point.

127.5 mL

c. $54.6\text{ mg} - 25\text{ mg} = 29.6\text{ mg}$

Because 25 has zero digits after the decimal point, round the answer to the nearest whole number.

30. mg

d. $2.35\text{ s} - 0.266\text{ s} = 2.084\text{ s}$

Because 2.35 has two digits after the decimal point, round the answer to two digits after the decimal point.

2.08 s

1.7 To write a number in scientific notation:

[1] Move the decimal point to give a number between 1 and 10.

[2] Multiply the result by 10^x , where x is the number of places the decimal point was moved.

a. $93,200 = 9.32 \times 10^4$

The decimal point was moved four places to the left.

c. $6,780,000 = 6.78 \times 10^6$

The decimal point was moved six places to the left.

e. $4,520,000,000,000 = 4.52 \times 10^{12}$

The decimal point was moved 12 places to the left.

b. $0.000\ 725 = 7.25 \times 10^{-4}$

The decimal point was moved four places to the right.

d. $0.000\ 030 = 3.0 \times 10^{-5}$

The decimal point was moved five places to the right.

f. $0.000\ 000\ 000\ 028 = 2.8 \times 10^{-11}$

The decimal point was moved 11 places to the right.

1.8 The exponent in 10^x tells how many places to move the decimal point in the coefficient to generate a standard number. The decimal point goes to the right when x is positive and to the left when x is negative.

a. $6.5 \times 10^3 = 6,500$

The decimal point was moved three places to the right.

b. $3.26 \times 10^{-5} = 0.000\ 032\ 6$

The decimal point was moved five places to the left.

c. $3.780 \times 10^{-2} = 0.037\ 80$

The decimal point was moved two places to the left.

d. $1.04 \times 10^8 = 104,000,000$

The decimal point was moved eight places to the right.

e. $2.221 \times 10^6 = 2,221,000$

The decimal point was moved six places to the right.

f. $4.5 \times 10^{-10} =$

0.000 000 000 45

The decimal point was moved 10 places to the left.

1.9 Use the equalities in Tables 1.3 and 1.4 to write a fraction that shows the relationship between the two units.

a. $\frac{0.621\ \text{mi}}{1\ \text{km}} \quad \frac{1\ \text{km}}{0.621\ \text{mi}}$

c. $\frac{454\ \text{g}}{1\ \text{lb}} \quad \frac{1\ \text{lb}}{454\ \text{g}}$

b. $\frac{1000\ \text{mm}}{1\ \text{m}} \quad \frac{1\ \text{m}}{1000\ \text{mm}}$

d. $\frac{1000\ \mu\text{g}}{1\ \text{mg}} \quad \frac{1\ \text{mg}}{1000\ \mu\text{g}}$

1.10 Use conversion factors to solve the problems.

a. $25\ \cancel{\text{L}} \times \frac{10\ \text{dL}}{1\ \cancel{\text{L}}} = 250\ \text{dL}$

b. $40.0\ \cancel{\text{oz}} \times \frac{28.3\ \text{g}}{1\ \cancel{\text{oz}}} = 1,132\ \text{g} = 1,130\ \text{g}$ rounded to 3 significant figures

c. $32\ \cancel{\text{in.}} \times \frac{2.54\ \text{cm}}{1\ \cancel{\text{in.}}} = 81.28\ \text{cm} = 81\ \text{cm}$ rounded to 2 significant figures

d. $10\ \cancel{\text{cm}} \times \frac{10\ \text{mm}}{1\ \cancel{\text{cm}}} = 100\ \text{mm}$

1.11 Use conversion factors to solve the problems.

a. $6,250\ \cancel{\text{ft}} \times \frac{1\ \cancel{\text{mi}}}{5,280\ \cancel{\text{ft}}} \times \frac{1\ \text{km}}{0.621\ \cancel{\text{mi}}} = 1.91\ \text{km}$
 Feet cancel. Miles cancel. Kilometers do not cancel.

b. $3\ \cancel{\text{cups}} \times \frac{1\ \cancel{\text{qt}}}{4\ \cancel{\text{cups}}} \times \frac{1\ \text{L}}{1.06\ \cancel{\text{qt}}} = 0.7\ \text{L}$
 Cups cancel. Quarts cancel. Liters do not cancel.

$$c. \quad 4.5 \cancel{\text{ft}} \times \frac{12 \cancel{\mu\text{in.}}}{1 \cancel{\text{ft}}} \times \frac{2.54 \text{ cm}}{1 \cancel{\mu\text{in.}}} = 140 \text{ cm}$$

Centimeters do not cancel.
Feet cancel.
Inches cancel.

1.12

$$0.100 \cancel{\text{mg}} \times \frac{1000 \cancel{\mu\text{g}}}{1 \cancel{\text{mg}}} \times \frac{1 \text{ tablet}}{25 \cancel{\mu\text{g}}} = 4 \text{ tablets}$$

1.13 Use the conversion factors: 1 teaspoon = 5 mL and 80. mg acetaminophen/2.5 mL of Tylenol.

$$a. \quad 2.5 \cancel{\text{tsp}} \times \frac{5 \text{ mL}}{1 \cancel{\text{tsp}}} = 12.5 \text{ mL rounded to } 13 \text{ mL (2 significant figures)}$$

$$b. \quad 13 \cancel{\text{mL}} \times \frac{80. \text{ mg}}{2.5 \cancel{\text{mL}}} = 416 \text{ mg rounded to } 420 \text{ mg (2 significant figures)}$$

1.14 Convert from T_C to T_F and T_K using the formulas listed in Section 1.9.

$$T_F = 1.8(T_C) + 32 \qquad T_K = T_C + 273$$

$$= 1.8(28.5) + 32 = 83.3 \text{ }^\circ\text{F} \qquad = 28.5 + 273 = 302 \text{ K}$$

- 1.15
- Because the densities of **A** and **B** are the same and there is a larger volume of **B**, the mass of **B** is greater than the mass of **A**.
 - The density of **A** is twice the density of **B**, but there is three times as much volume of **B** as **A**, so the mass of **B** is greater than the mass of **A**.
 - The density of **B** is greater than the density of **A** and there is a larger volume of **B**, so the mass of **B** is greater than the mass of **A**.

1.16 To convert volume (mL) to mass (g), multiply the volume by the density (g/mL).

$$10.0 \cancel{\text{mL}} \times \frac{0.713 \text{ g}}{1 \cancel{\text{mL}}} = 7.13 \text{ g}$$

Milliliters cancel.

Solutions to In-Chapter Problems

1.1 Naturally occurring: ice, blood. Synthetic: gloves, mask, plastic syringe, stainless steel needle.

1.2 *Physical properties* can be observed or measured without changing the composition of the material (a and d). *Chemical properties* determine how a substance can be converted to another substance by chemical reactions (b, c, and e).

- 1.3 This represents a chemical change because the “particles” on the left are different from the particles on the right. For example, on the left side there are particles containing only two red balls, whereas on the right there are none of these.
- 1.4 a. **B** and **C** illustrate pure elements.
b. **A** is a mixture of a compound and an element.
c. **D** is a mixture of two compounds.
- 1.5 A *pure substance* is composed of a single substance and has a constant composition regardless of the sample size (d). A *mixture* is composed of more than one component (a, b, and c). The composition of a mixture can vary depending on the sample.
- 1.6 An *element* is a pure substance that cannot be broken down into simpler substances by a chemical reaction (a). A *compound* is a pure substance formed by combining two or more elements together (b, c, and d).
- 1.7 One nanometer = 0.000 000 001 m (one billionth of a meter); therefore, 1 m = 1,000,000,000 nm.
- 1.8 a. 0.000 001 g = one microgram (1 μ g)
b. 1,000,000,000 m = one gigameter (1 Gm)
c. 0.000 000 001 s = one nanosecond (1 ns)
d. 0.01 g = one centigram (1 cg)
- 1.9 Use Table 1.2 to determine which quantity is larger.
- | | |
|--------------|---------|
| a. 3 cL | c. 5 km |
| b. 1 μ g | d. 2 mL |
- 1.10 A zero is significant only if it occurs between two nonzero digits, or at the end of a number with a decimal point.
- | | | | |
|--|--------------------------------------|---|--|
| a. $\overbrace{0.003}^{\text{No}}$ $\overbrace{04}^{\text{Yes}}$ | b. $\overbrace{26,045}^{\text{Yes}}$ | c. $\overbrace{1,000,034}^{\text{Yes}}$ | d. $\overbrace{0.304}^{\text{No}}$ $\overbrace{00}^{\text{Yes}}$ |
|--|--------------------------------------|---|--|
- 1.11 a. 5.43^{05} standard form: 543,000; scientific notation: 5.43×10^5
b. 4.43^{-06} standard form: 0.000 004 43; scientific notation: 4.43×10^{-6}
c. 7.4^{04} standard form: 74,000; scientific notation: 7.4×10^4

1.12

$$9 \text{ lb} \times \frac{16 \text{ oz}}{1 \text{ lb}} = 144 \text{ oz} \quad 144 \text{ oz} + 8 \text{ oz} = 152 \text{ oz}$$

$$152 \cancel{\text{oz}} \times \frac{28.3 \cancel{\text{g}}}{1 \cancel{\text{oz}}} \times \frac{1 \text{ kg}}{1000 \cancel{\text{g}}} = 4.3 \text{ kg}$$

1.13

$$0.46 \cancel{\text{mL}} \times \frac{1 \cancel{\text{L}}}{1000 \cancel{\text{mL}}} \times \frac{1,000,000 \mu\text{L}}{1 \cancel{\text{L}}} = 460 \mu\text{L}$$

1.14

$$160 \cancel{\text{mg}} \times \frac{5 \text{ mL}}{100 \cancel{\text{mg}}} = 8 \text{ mL Children's Motrin}$$

1.15

$$\begin{aligned} \text{a. } T_F &= 1.8(T_C) + 32 \\ &= 1.8(20.) + 32 = 68 \text{ }^\circ\text{F} \end{aligned}$$

$$\begin{aligned} \text{c. } T_C &= T_K - 273 \\ &= 298 - 273 = 25 \text{ }^\circ\text{C} \\ T_F &= 1.8(T_C) + 32 \\ &= 1.8(25) + 32 = 77 \text{ }^\circ\text{F} \end{aligned}$$

$$\begin{aligned} \text{b. } T_C &= \frac{T_F - 32}{1.8} \\ &= \frac{150. - 32}{1.8} = 66 \text{ }^\circ\text{C} \end{aligned}$$

$$\begin{aligned} \text{d. } T_K &= T_C + 273 \\ &= 75 + 273 = 348 \text{ K} \end{aligned}$$

1.16 Convert pounds of lead to grams. Then use the density of lead (11.3 g/cc) to determine the volume.

$$5 \text{ weights} \times \frac{2.0 \text{ lb}}{1 \text{ weight}} \times \frac{454 \text{ g}}{1 \cancel{\text{lb}}} \times \frac{1 \text{ cc}}{11.3 \cancel{\text{g}}} = \frac{402 \text{ cc}}{4.0 \times 10^2 \text{ cc lead}}$$

1.17

$$10.5 \cancel{\text{kg}} \times \frac{1000 \cancel{\text{g}}}{1 \cancel{\text{kg}}} \times \frac{1 \cancel{\text{mL}}}{1.38 \cancel{\text{g}}} \times \frac{1 \text{ L}}{1000 \cancel{\text{mL}}} = 7.61 \text{ L}$$

1.18

$$\text{a. specific gravity} = \frac{\text{density of a substance (g/mL)}}{\text{density of water (g/mL)}} = \frac{0.80 \text{ g/mL}}{1 \text{ g/mL}} = 0.80$$

$$\text{b. } 2.3 = \frac{\text{density of a substance (g/mL)}}{1 \text{ g/mL}} \quad \text{density} = 2.3 \text{ g/mL}$$

Solutions to End-of-Chapter Problems

- 1.19 Representation (a) is a pure element because the particle consists of a gray sphere. Representation (b) is a pure compound because each particle contains four gray spheres and one black sphere. Representation (c) is a mixture, because some of the particles contain two gray spheres, whereas others contain four gray spheres and one black sphere. Representation (d) is a mixture, because some of the particles are gray spheres, and some are blue.
- 1.20 Molecular art for an element shows spheres of one color only.
Elements: two blue spheres joined, two red spheres joined
Compounds: black sphere joined to two red spheres, red sphere joined to two gray spheres
- 1.21 a. **D** represents a mixture of two elements.
b. **B** is a mixture of a compound and an element.
- 1.22 a. physical change b. chemical change c. chemical change
- 1.23 A *chemical change* converts one substance to another substance by a chemical reaction (b). A *physical change* can be observed or measured without changing the composition of the material (a and c).
- 1.24 a. physical change b. physical change c. chemical change
- 1.25 This is a physical change, because the compound CO_2 is unchanged in this transition. The same “particles” exist at the beginning and end of the process.
- 1.26 A chemical change has occurred. Molecules of H_2 and N_2 have been converted into molecules of NH_3 (ammonia).
- 1.27 a, b: The temperature on the Fahrenheit thermometer is 76.5°F , which has three significant figures.
- c. $T_C = \frac{T_F - 32}{1.8} = \frac{76.5 - 32}{1.8} = 24.7^\circ\text{C}$
- 1.28 a. The length of the crayon is 4.5 cm. b. This value contains two significant figures.
- c. $4.5 \text{ cm} \times \frac{1 \text{ m}}{100 \text{ cm}} = 4.5 \times 10^{-2} \text{ m}$

- 1.29 a. 10 cloves of garlic: exact; two tablespoons of oil: inexact
b. five puppies: exact; 10 lb: inexact
- 1.30 a. 4 bicycles: exact number; 250 mi: inexact number
b. 4 cm: inexact number; 12 stitches: exact number
- 1.31 Compare the measurements using Table 1.2. (< means *less than*; > means *greater than*.)
- a. 5 mL < 5 dL b. 10 mg > 10 μg c. 5 cm > 5 mm d. 10 Ms > 10 ms
- 1.32 Compare the measurements using Table 1.2. (< means *less than*; > means *greater than*.)
- a. 10 km > 10 m b. 10 L > 10 mL c. 10 g > 10 μg d. 10 cm > 10 mm
- 1.33 All nonzero digits are significant. A zero is significant only if it occurs between two nonzero digits, or at the end of a number with a decimal point. The significant figures are in **bold**.
- | | | |
|--------------------------------------|---------------------------------------|--|
| a. 16.00 4 significant figures | d. 1,600,000 2 significant figures | g. 1.060 × 10 ¹⁰ 4 significant figures |
| b. 160 2 significant figures | e. 1.06 3 significant figures | h. 1.6 × 10 ⁻⁶ 2 significant figures |
| c. 0.001 60 3 significant figures | f. 0.1600 4 significant figures | |
- 1.34 All nonzero digits are significant. A zero is significant only if it occurs between two nonzero digits, or at the end of a number with a decimal point. The significant figures are in **bold**.
- | | | |
|--------------------------------------|------------------------------------|---|
| a. 160. 3 significant figures | d. 1.60 3 significant figures | g. 1.600 × 10 ⁻¹⁰ 4 significant figures |
| b. 160.0 4 significant figures | e. 1,600. 4 significant figures | h. 1.6 × 10 ⁶ 2 significant figures |
| c. 0.000 16 2 significant figures | f. 1.060 4 significant figures | |
- 1.35 When the number to be rounded off is 4 or fewer, it and all other digits to the right are dropped. When the number is 5 or greater, 1 is added to the digit to its left.
- | | | |
|--------------------------|-----------------------------|------------------|
| a. 25,401 = 25,400 | c. 0.001 265 982 = 0.001 27 | e. 195.371 = 195 |
| b. 1,248,486 = 1,250,000 | d. 0.123 456 = 0.123 | f. 196.814 = 197 |

- 1.36 When the number to be rounded off is 4 or fewer, it and all other digits to the right are dropped. When the number is 5 or greater, 1 is added to the digit to its left.

a. $25,401 = 2.540 \times 10^4$ c. $0.001\ 265\ 982 = 0.001\ 266$ e. $195.371 = 195.4$
b. $1,248,486 = 1,248,000$ d. $0.123\ 456 = 0.123\ 5$ f. $196.814 = 196.8$

- 1.37 The answers in problems with multiplication and division must have the same number of significant figures as the original number with the fewest significant figures. The answers in problems with addition and subtraction must have the same number of digits after the decimal point as the original number with the fewest digits after the decimal point.

a. $53.6 \times 0.41 = 21.976$
Because 0.41 has two significant figures, round the answer to two significant figures.
22

c. $65.2 \div 12 = 5.43333$
Because 12 has two significant figures, round the answer to two significant figures.
5.4

e. $694.2 \times 0.2 = 138.84$
Because 0.2 has one significant figure, round the answer to one significant figure.
100

b. $25.825 - 3.86 = 21.965$
Because 3.86 has two digits after the decimal point, round the answer to two digits after the decimal point.
21.97

d. $41.0 + 9.135 = 50.135$
Because 41.0 has one digit after the decimal point, round the answer to one digit after the decimal point.
50.1

f. $1,045 - 1.26 = 1,043.74$
Because 1,045 has no digits after the decimal point, round the answer to the closest whole number.
1,044

- 1.38 The answers in problems with multiplication and division must have the same number of significant figures as the original number with the fewest significant figures. The answers in problems with addition and subtraction must have the same number of digits after the decimal point as the original number with the fewest digits after the decimal point.

a. $49,682 \times 0.80 = 39,745.60$
Since 0.80 has two significant figures, round the answer to two significant figures.
 4.0×10^4

c. $1,000 \div 2.34 = 427.35$
Since 1,000 has one significant figure, round the answer to one significant figure.
400

e. $25,000 \div 0.4356 = 57,392.10$
Since 25,000 has two significant figures, round the answer to two significant figures.
57,000

b. $66.815 + 2.82 = 69.635$
Since 2.82 has two digits after the decimal point, round the answer to two digits after the decimal point.
69.64

d. $21 - 0.88 = 20.12$
Since 21 has no digits after the decimal point, round the answer to the closest whole number.
20.

f. $21.5381 + 26.55 = 48.0881$
Since 26.55 has two digits after the decimal point, round the answer to two digits after the decimal point.
48.09

1.39 To write a number in scientific notation:

[1] Move the decimal point to give a number between 1 and 10.

[2] Multiply the result by 10^x , where x is the number of places the decimal point was moved.

a. $1,234 \text{ g} = 1.234 \times 10^3 \text{ g}$

The decimal point was moved three places to the left.

b. $0.000\ 016\ 2 \text{ m} = 1.62 \times 10^{-5} \text{ m}$

The decimal point was moved five places to the right.

c. $5,244,000 \text{ L} = 5.244 \times 10^6 \text{ L}$

The decimal point was moved six places to the left.

d. $0.005\ 62 \text{ g} = 5.62 \times 10^{-3} \text{ g}$

The decimal point was moved three places to the right.

e. $44,000 \text{ km} = 4.4 \times 10^4 \text{ km}$

The decimal point was moved four places to the left.

1.40 To write a number in scientific notation:

[1] Move the decimal point to give a number between 1 and 10.

[2] Multiply the result by 10^x , where x is the number of places the decimal point was moved.

a. $0.001\ 25 \text{ m} = 1.25 \times 10^{-3} \text{ m}$

The decimal point was moved three places to the right.

b. $8,100,000,000 \text{ lb} = 8.1 \times 10^9 \text{ lb}$

The decimal point was moved nine places to the left.

c. $54,235.6 \text{ m} = 5.42356 \times 10^4 \text{ m}$

The decimal point was moved four places to the left.

d. $0.000\ 001\ 899 \text{ L} = 1.899 \times 10^{-6} \text{ L}$

The decimal point was moved six places to the right.

e. $4,440 \text{ s} = 4.44 \times 10^3 \text{ s}$

The decimal point was moved three places to the left.

1.41 The exponent in 10^x tells how many places to move the decimal point in the coefficient to generate a standard number. The decimal point goes to the right when x is positive and to the left when x is negative.

a. $3.4 \times 10^8 = 340,000,000$

The decimal point was moved eight places to the right.

b. $5.822 \times 10^{-5} = 0.000\ 058\ 22$

The decimal point was moved five places to the left.

c. $3 \times 10^2 = 300$

The decimal point was moved two places to the right.

d. $6.86 \times 10^{-8} = 0.000\ 000\ 068\ 6$

The decimal point was moved eight places to the left.

1.42 The exponent in 10^x tells how many places to move the decimal point in the coefficient to generate a standard number. The decimal point goes to the right when x is positive and to the left when x is negative.

a. $4.02 \times 10^{10} = 40,200,000,000$

The decimal point was moved 10 places to the right.

b. $2.46 \times 10^{-3} = 0.002\ 46$

The decimal point was moved three places to the left.

c. $6.86 \times 10^9 = 6,860,000,000$

The decimal point was moved nine places to the right.

d. $1.00 \times 10^{-7} = 0.000\ 000\ 100$

The decimal point was moved seven places to the left.

1.43 Compare the two numbers. The number in **bold** is larger.

a. 4.44×10^3 or 4.8×10^2

b. 5.6×10^{-6} or 5.6×10^{-5}

c. 1.3×10^8 or 52,300,000

d. 9.8×10^{-4} or 0.000 089

1.44 a. $7 \times 10^4 < 5.06 \times 10^6 < 2.5 \times 10^8$

b. $8.6 \times 10^{-6} < 2.5 \times 10^{-4} < 6.3 \times 10^{-2}$

1.45 Write the number in scientific notation.

a. 0.000 400 g of folate = 4.00×10^{-4} g

The decimal point was moved four places to the right.

b. 0.002 g of copper = 2×10^{-3} g

The decimal point was moved three places to the right.

c. 0.000 080 g of vitamin K = 8.0×10^{-5} g

The decimal point was moved five places to the right.

d. 3,400 mg of chloride = 3.4×10^3 mg

The decimal point was moved three places to the left.

1.46 Use conversion factors to solve the problem and write the answer in scientific notation.

a. $0.40\ \mu\text{m} \times \frac{1\ \text{m}}{1 \times 10^6\ \mu\text{m}} = 4.0 \times 10^{-7}\ \text{m}$

b. $4.0 \times 10^{-7}\ \mu\text{m} \times \frac{100\ \text{cm}}{1\ \mu\text{m}} \times \frac{1\ \text{in}}{2.54\ \text{cm}} = 1.6 \times 10^{-5}\ \text{in}$

1.47 The scale shows the individual has a mass of 115 lb.

$115\ \text{lb} \times \frac{1\ \text{kg}}{2.20\ \text{lb}} = 52.3\ \text{kg}$

1.48 The syringe contains 1.4 mL of liquid.

a. 1.4 mL b. $1.4 \cancel{\text{mL}} \times \frac{1 \text{ L}}{1000 \cancel{\text{mL}}} = 0.0014 \text{ L} = 1.4 \times 10^{-3} \text{ L}$

1.49

a. $60.6 \cancel{\text{kg}} \times \frac{2.20 \text{ lb}}{1 \cancel{\text{kg}}} = 133 \text{ lb}$

b. $67 \cancel{\text{in}} \times \frac{2.54 \text{ cm}}{1 \cancel{\text{in}}} = 170 \text{ cm}$

1.50 Use conversion factors to solve the problems.

a. $3.5 \text{ tablespoons} \times \frac{15 \text{ mL}}{1 \text{ tablespoon}} = 53 \text{ mL}$

b. $\frac{53 \text{ mL}}{\text{dosage}} \times \frac{4 \text{ dosages}}{\text{day}} \times \frac{7 \text{ days}}{1 \text{ week}} \times 1 \text{ week} = 1.5 \times 10^3 \text{ mL}$

c. $1.5 \times 10^3 \cancel{\text{mL}} \times \frac{1 \text{ liter}}{1000 \cancel{\text{mL}}} = 1.5 \text{ L}$

1.51 Use conversion factors to solve the problem.

$$6.0 \cancel{\text{oz}} \times \frac{28.4 \cancel{\text{g}} \text{ fish}}{1 \cancel{\text{oz}} \text{ fish}} \times \frac{354 \text{ mg Hg}}{10^6 \cancel{\text{g}} \text{ fish}} = \frac{60. \times 10^3}{10^6} = 60. \times 10^{-3} \\ = 0.060 \text{ mg Hg}$$

1.52 Use conversion factors to solve the problem.

$$7 \cancel{\text{days}} \times \frac{875 \cancel{\text{mg}}}{1 \cancel{\text{day}}} \times \frac{1 \text{ gram}}{1000 \cancel{\text{mg}}} = 6.13 \text{ g K}$$

1.53 Use conversion factors to solve the problems.

a. $50. \cancel{\text{in}} \times \frac{2.54 \text{ cm}}{1 \cancel{\text{in}}} = 127 \text{ cm} = \mathbf{130 \text{ cm}}$ rounded to two significant figures

b. $3.0 \text{ pints} \times \frac{1 \cancel{\text{qt}}}{2 \cancel{\text{pints}}} \times \frac{1 \text{ L}}{1.06 \cancel{\text{qt}}} = 1.415 \text{ L} = \mathbf{1.4 \text{ L}}$ rounded to two significant figures

c. $T_F = 1.8(T_C) + 32$
 $= 1.8(37.7) + 32 = 99.9 \text{ }^\circ\text{F}$


1.60

Density of honey > water > gasoline

$$25 \cancel{\text{g}} \text{ water} \times \frac{1 \text{ mL}}{1.00 \cancel{\text{g}}} = 25 \text{ mL water}$$

$$15.0 \cancel{\text{g}} \text{ gasoline} \times \frac{1 \text{ mL}}{0.75 \cancel{\text{g}}} = 20. \text{ mL gasoline}$$

$$42.6 \cancel{\text{g}} \text{ honey} \times \frac{1 \text{ mL}}{1.42 \cancel{\text{g}}} = 30.0 \text{ mL honey}$$



- 1.61 a. Hexane is *less* dense than water, so 50 mL of hexane will be above the 100 mL of water.
b. Dichloromethane is *more* dense than water, so the 100 mL of water will be on top of the 50 mL of dichloromethane.

- 1.62 a. The density of the liquid in beaker **A** is less than 2.0 g/cc.
b. The density of the liquid in beaker **B** is greater than 0.90 g/cc.

1.63

$$\frac{122 \text{ g}}{121 \text{ mL}} = 1.01 \text{ g/mL}$$

- 1.64 Calculate the volume of water using the density of water; use that volume to calculate the density of the saline.

$$24.5 \cancel{\text{g}} \times \frac{1 \text{ mL}}{1.00 \cancel{\text{g}}} = 24.5 \text{ mL} \quad \frac{25.6 \text{ g}}{24.5 \text{ mL}} = 1.04 \text{ g/mL}$$

1.65

$$1 \cancel{\text{qt}} \times \frac{946 \cancel{\text{mL}}}{1 \cancel{\text{qt}}} \times \frac{1.03 \cancel{\text{g}}}{1 \cancel{\text{mL}}} \times \frac{1 \text{ kg}}{1000 \cancel{\text{g}}} = 0.974 \text{ 38 kg} = \mathbf{0.974 \text{ kg}}$$

1.66

$$\frac{10.1 \text{ g}}{4.1 \text{ mL}} = 2.5 \text{ g/mL}$$

1.67 The density of a substance determines whether it floats or sinks in a liquid. The less dense liquid is the upper layer. The density of water is 1.0 g/mL.

- | | |
|--|------------------------------------|
| a. heptane (0.684 g/mL < 1.0 g/mL) | c. water (1.0 g/mL < 1.49 g/mL) |
| b. olive oil (0.92 g/mL < 1.0 g/mL) | d. water (1.0 g/mL < 1.59 g/mL) |

1.68

$$\text{a. specific gravity} = \frac{\text{density of mercury (g/mL)}}{\text{density of water (g/mL)}} = \frac{13.6 \text{ g/mL}}{1 \text{ g/mL}} = 13.6$$

$$\text{b. } 0.789 = \frac{\text{density of ethanol (g/mL)}}{1 \text{ g/mL}} \quad \text{density} = 0.789 \text{ g/mL}$$

1.69 Use conversion factors to solve the problems.

$$\text{a. } \frac{186 \text{ mg}}{\text{dL}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 0.186 \text{ g/dL} \quad \text{b. } \frac{186 \text{ mg}}{\text{dL}} \times \frac{10 \text{ dL}}{1 \text{ L}} = 1,860 \text{ mg/L}$$

1.70

$$\text{a. } 88.0 \text{ } \cancel{\mu\text{g}} \times \frac{1 \cancel{\text{g}}}{1,000,000 \cancel{\mu\text{g}}} \times \frac{1000 \text{ mg}}{1 \cancel{\text{g}}} = 8.80 \times 10^{-2} \text{ mg}$$

$$\text{b. } \frac{8.80 \times 10^{-2} \cancel{\text{mg}}}{\text{day}} \times \frac{1 \text{ g}}{1000 \cancel{\text{mg}}} \times 30 \text{ days} = 2.64 \times 10^{-3} \text{ g}$$

1.71

$$1.5 \cancel{\text{g}} \times \frac{1000 \cancel{\text{mg}}}{1 \cancel{\text{g}}} \times \frac{1 \text{ tablet}}{500 \cancel{\text{mg}}} = 3 \text{ tablets}$$

1.72

$$1.8 \cancel{\text{L}} \times \frac{1000 \cancel{\text{mL}}}{1 \cancel{\text{L}}} \times \frac{1.05 \cancel{\text{g}}}{1 \cancel{\text{mL}}} \times \frac{1 \text{ kg}}{1000 \cancel{\text{g}}} = 1.9 \text{ kg}$$

$$70.7 \text{ kg} + 1.9 \text{ kg} = 72.6 \text{ kg}; 72.6 \text{ kg} - 69.3 \text{ kg} = \mathbf{3.3 \text{ kg sweat lost}}$$

$$3.3 \cancel{\text{kg}} \times \frac{2.20 \cancel{\text{lb}}}{1 \cancel{\text{kg}}} = \mathbf{7.3 \text{ lb}}$$

1.73

$$232 \cancel{\text{lb}} \times \frac{1 \cancel{\text{kg}}}{2.20 \cancel{\text{lb}}} \times \frac{25 \text{ kg body fat}}{100 \cancel{\text{kg}}} = 26 \text{ kg body fat}$$

1.74

$$13.0 \text{ oz} \times \frac{250 \text{ mg}}{1 \text{ oz}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 3.25 \text{ g of sodium}$$

3.25 g - 2.4 g = 0.9 g more sodium than recommended daily value

1.75

$$\text{a. } \frac{20 \text{ mL}}{1 \text{ dose}} \times \frac{\$10.00}{300. \text{ mL}} = \frac{\$0.67}{\text{dose}}$$

b. 2 tablespoons = 30. mL

$$\frac{30 \text{ mL}}{1 \text{ dose}} \times \frac{\$10.00}{300. \text{ mL}} = \frac{\$1.00}{\text{dose}}$$

1.76

$$\text{a. } 1.93 \text{ mg} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 1.93 \times 10^{-3} \text{ g} \quad 1.93 \times 10^{-3} \text{ g} \times \frac{1,000,000 \text{ } \mu\text{g}}{1 \text{ g}} = 1.93 \times 10^3 \text{ } \mu\text{g}$$

$$\text{b. } 20 \text{ cigarettes} \times \frac{1.93 \text{ mg}}{\text{cigarette}} = 38.6 \text{ mg} \quad 38.6 \text{ mg} > 21 \text{ mg}; \text{ the smoker will get less nicotine from the patch}$$

1.77

2 tablets x 325 mg/tablet = 650. mg

$$0.510 \text{ kg} \times \frac{1000 \text{ g}}{1 \text{ kg}} \times \frac{1000 \text{ mg}}{1 \text{ g}} \times \frac{1 \text{ dose}}{650. \text{ mg}} = 784.6 = 784 \text{ full doses}$$

1.78

$$\frac{4 \text{ times}}{\text{day}} \times \frac{2 \text{ tsp}}{\text{time}} \times \frac{5 \text{ mL}}{1 \text{ tsp}} \times \frac{400 \text{ mg Al(OH)}_3}{5 \text{ mL}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 3.2 \text{ g Al(OH)}_3$$

$$\frac{4 \text{ times}}{\text{day}} \times \frac{2 \text{ tsp}}{\text{time}} \times \frac{5 \text{ mL}}{1 \text{ tsp}} \times \frac{400 \text{ mg Mg(OH)}_2}{5 \text{ mL}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 3.2 \text{ g Mg(OH)}_2$$

$$\frac{4 \text{ times}}{\text{day}} \times \frac{2 \text{ tsp}}{\text{time}} \times \frac{5 \text{ mL}}{1 \text{ tsp}} \times \frac{40 \text{ mg simethicone}}{5 \text{ mL}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 0.32 \text{ g simethicone}$$

1.79

$$4 \text{ times} \times \frac{2.0 \text{ g}}{\text{time}} \times \frac{1000 \text{ mg}}{1 \text{ g}} \times \frac{1 \text{ tablet}}{500. \text{ mg}} = 16 \text{ tablets}$$

1.80

$$\text{a. } 20. \cancel{\text{min}} \times \frac{1 \cancel{\text{h}}}{60 \cancel{\text{min}}} \times \frac{150 \text{ mL}}{1 \cancel{\text{h}}} = 50. \text{ mL}$$

$$\text{b. } 90. \cancel{\text{mL}} \times \frac{1 \cancel{\text{h}}}{150 \cancel{\text{mL}}} \times \frac{60 \text{ min}}{1 \cancel{\text{h}}} = 36 \text{ min}$$

$$\text{c. } 600. \cancel{\text{mL}} \times \frac{1 \text{ h}}{150 \cancel{\text{mL}}} = 4.0 \text{ hours}$$

$$\text{d. } 2.0 \cancel{\text{g}} \times \frac{1000 \cancel{\text{mg}}}{1 \cancel{\text{g}}} \times \frac{\cancel{\text{mL}}}{90. \cancel{\text{mg}}} \times \frac{1 \cancel{\text{h}}}{150 \cancel{\text{mL}}} \times \frac{60 \text{ min}}{1 \cancel{\text{h}}} = 8.9 \text{ min}$$

1.81

$$\frac{2.0 \text{ mg}}{1 \cancel{\text{kg}}} \times \frac{1 \cancel{\text{kg}}}{2.20 \cancel{\text{lb}}} \times 110 \cancel{\text{lb}} = 1.0 \times 10^2 \text{ mg}$$

1.82

$$28 \cancel{\text{kg}} \times \frac{10 \cancel{\text{mg}}}{\cancel{\text{kg}} \text{ dose}} \times \frac{3 \text{ doses}}{\cancel{\text{day}}} \times 7 \cancel{\text{days}} \times \frac{1 \text{ g}}{1000 \cancel{\text{mg}}} = 5.9 \text{ g}$$

1.83 Convert mass and height to kg and m, respectively. Use the formula, $\text{BMI} = \text{kg}/\text{m}^2$, to solve the problem.

$$180 \cancel{\text{lb}} \times \frac{1 \text{ kg}}{2.20 \cancel{\text{lb}}} = 82 \text{ kg}$$

$$6 \text{ ft } 1 \text{ in.} = 73 \text{ in.}$$

$$73 \cancel{\text{in.}} \times \frac{1 \text{ m}}{39.4 \cancel{\text{in.}}} = 1.9 \text{ m}$$

$$\text{BMI} = \frac{\text{kg}}{\text{m}^2} = \frac{82}{(1.9)^2} = \frac{82}{3.61} = 23$$

The BMI is in the normal range.

1.84

$$\text{a. } 150. \cancel{\text{lb}} \times \frac{1 \text{ kg}}{2.20 \cancel{\text{lb}}} = 68.2 \text{ kg} \quad 3 \text{ tablets} \times \frac{200. \cancel{\text{mg}}}{1 \cancel{\text{tablet}}} = 600. \text{ mg}$$

$$\frac{600. \text{ mg}}{68.2 \text{ kg}} = 8.80 \text{ mg/kg}$$

$$\text{b. } 45 \cancel{\text{kg}} \times \frac{8.80 \cancel{\text{mg}}}{1 \cancel{\text{kg}}} = 4.0 \times 10^2 \text{ mg}$$

1.85

$$42 \text{ lb} \times \frac{1 \text{ kg}}{2.20 \text{ lb}} \times \frac{10 \text{ mg}}{1 \text{ kg}} \times \frac{1 \text{ tablet}}{80 \text{ mg}} = 2.4 = \mathbf{2 \text{ tablets}}$$

1.86

$$10.0 \text{ mg} + 4(80. \text{ mg}) = 330. \text{ mg} \quad \text{amount of dosage from tablets}$$

$$3.2 \text{ mg/kg} + 4(1.6 \text{ mg/kg}) = 9.6 \text{ mg/kg} \quad \text{amount of dosage from injection}$$

$$\text{a. } 40. \text{ kg} \times 9.6 \text{ mg/kg} = 3.8 \times 10^2 \text{ mg} \quad \text{tablet form gives higher dosage}$$

$$\text{b. } 100. \text{ kg} \times 9.6 \text{ mg/kg} = 9.6 \times 10^2 \text{ mg} \quad \text{injections give higher dosage}$$

1.87

$$1.5 \text{ tsp} \times \frac{5.0 \text{ mL}}{1 \text{ tsp}} \times \frac{100. \text{ mg}}{5 \text{ mL}} \times \frac{1 \text{ g}}{1000 \text{ mg}} = 0.15 \text{ g}$$

1.88

$$200 \text{ lb} \times \frac{1 \text{ kg}}{2.20 \text{ lb}} \times \frac{10 \text{ } \mu\text{g}}{1 \text{ kg}} \times \frac{1 \text{ mg}}{1000 \text{ } \mu\text{g}} = 0.909 \text{ mg} = \mathbf{0.9 \text{ mg}}$$